

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	1	clickb	US-PGPUB; USPAT; DERWENT	ADJ	ON	2005/03/07 15:39
L2	1094	Clc and chloride	US-PGPUB; USPAT; DERWENT	ADJ	ON	2005/03/07 15:40
L3	56	Clc and chloride.ab.	US-PGPUB; USPAT; DERWENT	ADJ	ON	2005/03/07 15:42
L4	114	CICK	US-PGPUB; USPAT; DERWENT	ADJ	ON	2005/03/07 15:42
L5	5	CICK and chloride	US-PGPUB; USPAT; DERWENT	ADJ	ON	2005/03/07 15:43
L6	0	CICK2b and chloride	US-PGPUB; USPAT; DERWENT	ADJ	ON	2005/03/07 15:43

10/687523

10 = 10-15-03

File 5:Biosis Previews(R) 1969-2005/Feb W4  
(c) 2005 BIOSIS

Set	Items	Description
Set	Items	Description
S1	4	CLCKB
S2	3	CHANNEL() KB
S3	274	AU='LANG FLORIAN' OR AU='LANG FLORIAN J W'
S4	67	AU='WALDEGGER SIEGFRIED' OR AU='WALDEGGER SIEGRIED'
S5	2	AU='LAMPERT ANGELIKA'
S6	51	S3 AND S4
S7	9	S6 AND CHLORIDE

? t s1/7/1-4

1/7/1  
DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0015143594 BIOSIS NO.: 200500050344  
Novel mutations of the chloride channel Kb gene in two Japanese patients clinically diagnosed as Bartter syndrome with hypocalciuria  
AUTHOR: Fukuyama Shigeru; Hiramatsu Misako; Akagi Motohiro; Higa Mutumi; Ohta Takao (Reprint)  
AUTHOR ADDRESS: Fac MedDept Pediat, Univ Ryukyus, 207 Uehara, Okinawa, 9030125, Japan\*\*Japan  
AUTHOR E-MAIL ADDRESS: tohta@med.u-ryukyu.ac.jp  
JOURNAL: Journal of Clinical Endocrinology & Metabolism 89 (11): p 5847-5850 November 2004 2004  
MEDIUM: print  
ISSN: 0021-972X (ISSN print)  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: Hypokalemic metabolic tubulopathy, such as in Bartter syndrome and Gitelman syndrome, is caused by the dysfunction of renal electrolyte transporters. Despite advances in molecular genetics with regard to hypokalemic metabolic tubulopathy, recent reports have suggested that the phenotype-genotype correlation is still confusing, especially in classic Bartter and Gitelman syndromes. We report here two Japanese patients who suffered from clinically diagnosed classic Bartter syndrome but who presented hypocalciuria. Hypocalciuria is generally believed to be a pathognomonic finding of NCCT malfunction. To better understand the genotype-phenotype correlation in these two cases, we screened four renal electrolyte transporter genes (Na-K-2Cl cotransporter (NKCC2), renal outer medullary K channel (ROMK), Cl channel Kb (\*\*\*ClCKb\*\*\*), and Na-Cl cotransporter (NCCT)) by the PCR direct sequencing method. We identified three ClC-Kb allelic variants, including two new mutations (L27R and W610X in patient 1 and a G to C substitution of a 3' splice-site of intron 2 and W610X in patient 2). We did not find any mutations in the other three genes. Our present data suggest that some ClC-Kb mutations may affect calcium handling in renal tubular cells.

1/7/2  
DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0013223243 BIOSIS NO.: 200100395082  
Bartter' syndrome type 3 (BS3): An unusual cause of nephrolithiasis  
AUTHOR: Colussi G (Reprint); De Ferrari M E (Reprint); Bettinelli A; Tedeschi S; Cesareo L; Civati G (Reprint)  
AUTHOR ADDRESS: Renal Unit, Riguarda-Ca' Granda Hospital, Milan, Italy\*\* Italy  
JOURNAL: Nephrology Dialysis Transplantation 16 (6): pA14 June, 2001 2001  
MEDIUM: print  
CONFERENCE/MEETING: Annual Congress of the European Renal Association and the European Dialysis and Transplant Association Vienna, Austria June

24-27, 2001; 20010624

SPONSOR: European Renal Association

European Dialysis and Transplant Association

ISSN: 0931-0509

DOCUMENT TYPE: Meeting; Meeting Abstract

RECORD TYPE: Citation

LANGUAGE: English

1/7/3

DIALOG(R)File 5:Biosis Previews(R)

(c) 2005 BIOSIS. All rts. reserv.

0011959401 BIOSIS NO.: 199900219061

Expression of CLCN voltage-gated chloride channel genes in human blood vessels

AUTHOR: Lamb Fred S (Reprint); Clayton Gerald H; Liu Bei-Xing; Smith Roderic L; Barna Thomas J; Schutte Brian C

AUTHOR ADDRESS: Department of Pediatrics, University of Iowa Hospitals, 5040C RCP, Iowa City, IA, 52242, USA\*\*USA

JOURNAL: Journal of Molecular and Cellular Cardiology 31 (3): p657-666 March, 1999 1999

MEDIUM: print

ISSN: 0022-2828

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: Expression of CLCN Voltage-gated Chloride Channel Genes in Human Blood Vessels. Journal of Molecular and Cellular Cardiology (1999) 31, 657-666. Chloride (Cl) ion channels play a critical role in the response of both vascular smooth muscle (VSM) and endothelial (ENDO) cells to agonist stimulation. In VSM, agonist-induced Cl currents produce membrane depolarization, resulting in calcium influx through voltage-sensitive channels. ENDO cells also activate Cl currents after either agonist application or perturbation of cell volume. Although some of these currents have been characterized biophysically, the genes involved have not been identified. The CLCN family of voltage-dependent Cl channel genes comprises nine members (CLCN1-7, Ka and Kb) which demonstrate quite diverse functional characteristics while sharing significant sequence homology. We used Northern-blot analysis to study the expression of these Cl channel genes in cultured human aortic and coronary VSM cells and in aortic ENDO cells. CLCN3 is by far the most abundant CLC channel mRNA in both VSM and ENDO cells. Lower levels of expression are seen for CLCN2, CLCN4, CLCN5 and CLCN6. Expression levels were similar in VSM and ENDO cells except for CLCN4 which was more highly expressed in ENDO cells. In situ hybridization was used to confirm the expression of CLCN3 in intact human fetal lung. CLCN3 message was seen in VSM and ENDO cells of both large and small pulmonary vessels, indicating that their detection by Northern blotting was not an artifact of cell culture. CLCN3 is also expressed in pulmonary epithelial and bronchial smooth muscle cells but not in chondrocytes or pulmonary interstitial cells. Recent studies suggest that CLCN3 may encode the swelling-induced Cl conductance. We used whole cell patch clamp recording to demonstrate swelling-induced Cl currents in these cultured VSM cells. This suggests that the CLCN3 protein is expressed; however, the functional role of this current in VSM remains to be determined.

1/7/4

DIALOG(R)File 5:Biosis Previews(R)

(c) 2005 BIOSIS. All rts. reserv.

0011858511 BIOSIS NO.: 199900118171

Gitelman syndrome due to mutation in the chloride channel \*\*\*CLCKB\*\*\*

AUTHOR: Jeck N; Konrad M; Reinalter S; Seyberth H W

AUTHOR ADDRESS: Philipps-Univ., Dep. Pediatr., Marburg, Germany\*\*Germany

JOURNAL: Kidney and Blood Pressure Research 21 (2-4): p180 1998 1998

MEDIUM: print

CONFERENCE/MEETING: Congress of Nephrology 1998 Joint Scientific Meeting of the Society Nephrology Erlangen, Germany September 19-22, 1998; 19980919

SPONSOR: The Society for Nephrology  
ISSN: 1420-4096  
DOCUMENT TYPE: Meeting; Meeting Abstract; Meeting Poster  
RECORD TYPE: Citation  
LANGUAGE: English

? t s2/7/1-3

2/7/1  
DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0015143594 BIOSIS NO.: 200500050344  
Novel mutations of the chloride ~~channel~~ ~~gene~~ ~~in two Japanese~~  
patients clinically diagnosed as Bartter syndrome with hypocalciuria  
AUTHOR: Fukuyama Shigeru; Hiramatsu Misako; Akagi Motohiro; Higa Mutumi;  
Ohta Takao (Reprint)  
AUTHOR ADDRESS: Fac MedDept Pediat, Univ Ryukyus, 207 Uehara, Okinawa,  
9030125, Japan\*\*Japan  
AUTHOR E-MAIL ADDRESS: tohta@med.u-ryukyu.ac.jp  
JOURNAL: Journal of Clinical Endocrinology & Metabolism 89 (11): p  
5847-5850 November 2004 2004  
MEDIUM: print  
ISSN: 0021-972X (ISSN print)  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: Hypokalemic metabolic tubulopathy, such as in Bartter syndrome  
and Gitelman syndrome, is caused by the dysfunction of renal electrolyte  
transporters. Despite advances in molecular genetics with regard to  
hypokalemic metabolic tubulopathy, recent reports have suggested that the  
phenotype-genotype correlation is still confusing, especially in classic  
Bartter and Gitelman syndromes. We report here two Japanese patients who  
suffered from clinically diagnosed classic Bartter syndrome but who  
presented hypocalciuria. Hypocalciuria is generally believed to be a  
pathognomonic finding of NCCT malfunction. To better understand the  
genotype-phenotype correlation in these two cases, we screened four renal  
electrolyte transporter genes (Na-K-2Cl cotransporter (NKCC2), renal  
outer medullary K channel (ROMK), Cl ~~channel~~ ~~gene~~ (ClCKb), and  
Na-Cl cotransporter (NCCT)) by the PCR direct sequencing method. We  
identified three ClC-Kb allelic variants, including two new mutations  
(L27R and W610X in patient 1 and a G to C substitution of a 3' splice  
site of intron 2 and W610X in patient 2). We did not find any mutations  
in the other three genes. Our present data suggest that some ClC-Kb  
mutations may affect calcium handling in renal tubular cells.

2/7/2  
DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0014478721 BIOSIS NO.: 200300447440  
Analysis of renal tubular electrolyte transporter genes in seven patients  
with hypokalemic metabolic alkalosis.  
AUTHOR: Fukuyama Shigeru; Okudaira Shoko; Yamazato Syosin; Yamazato  
Masahiro; Ohta Takao (Reprint)  
AUTHOR ADDRESS: Department of Pediatrics, Faculty of Medicine, University  
of the Ryukyus, 207 Uehara, Nishihara, Okinawa, 903-0125, Japan\*\*Japan  
AUTHOR E-MAIL ADDRESS: tohta@med.u-ryukyu.ac.jp  
JOURNAL: Kidney International 64 (3): p808-816 September 2003 2003  
MEDIUM: print  
ISSN: 0085-2538 (ISSN print)  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: Background: Disorders that manifest hypokalemic metabolic  
alkalosis, such as Bartter's syndrome and Gitelman's syndrome, are caused  
by the malfunction of renal tubular electrolyte transporters. Bartter's

syndrome may be linked to dysfunction of Na-K-2Cl cotransporter (NKCC2), renal outer medullary K channel (ROMK), or Cl ~~channel~~ ~~Kb~~ (ClC-Kb), while Gitelman's syndrome may be linked to Na-Cl cotransporter (NCCT) dysfunction. However, previous genetic analyses in these syndromes have included many heterozygotes for each gene and there has been no further analysis of other genes. Thus, to clarify the interaction of these transporter genes, in the present study we investigated all 4 transporter genes in 7 patients with hypokalemic metabolic alkalosis. Methods: Seven patients from 5 families (patients A-G) were collected, and a mutation analysis of the 4 renal electrolyte transporter genes was performed by direct sequencing. Results: We identified 12 mutations in these 7 patients. Three mutations (del245Y in NKCC2, R1009X in NCCT, V524I in ClC-Kb) have not been reported previously. In NKCC2 gene screening, patient A was homozygous for del245Y. In ClC-Kb gene screening, L27R was detected in patients B, D, and E. V524I was detected in patient C. Both T562M and E578K were observed in patients B and E. In NCCT gene screening, patients B-G shared a common novel mutation, R1009X, and patients D, E, F, and G carried this mutation in both alleles. Patients B and C carried R1009X in one allele, and a 6-amino acid insertion in exon 6 and L849H in another allele, respectively. The 4 other mutations did not result in any amino acid exchange. Despite the NCCT gene mutation, patients C and E showed normomagnesemia. Conclusion: Our findings demonstrate that in Bartter's and Gitelman's syndromes, it may not be uncommon to see mutations in several causative transporter genes.

2/7/3

DIALOG(R)File 5: Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0008926577 BIOSIS NO.: 199396090993

Effect of KB-2796, a new diphenylpiperazine calcium antagonist, on voltage-dependent calcium currents and oxidative metabolism in dissociated mammalian CNS neurons

AUTHOR: Akaike Norio (Reprint); Ishibashi Hitoshi; Hara Hideaki; Oyama Yasuo; Ueha Toshiko

AUTHOR ADDRESS: Dep. Neurophysiol., Tohoku Univ. Sch. Med., Sendai 980, Japan\*\*Japan

JOURNAL: Brain Research 619 (1-2): p263-270 1993

ISSN: 0006-8993

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: The effects of KB-2796,

1-(bis(4-fluorophenyl)methyl)-4-(2,3,4-trimethoxybenzyl)piperazine-2HCl, on the low- and high-voltage activated Ca-2+ currents (LVA and HVA I-Ca, respectively) and on oxidative metabolism were studied in neurons freshly dissociated from rat brain. KB-2796 reduced the peak amplitude of LVA I-Ca in a concentration-dependent manner with a threshold concentration of 10<sup>-7</sup> M when the LVA I-Ca was elicited every 30 s in the external solution with 10 mM Ca-2+. The concentration for half-maximum inhibition (IC-50) was 1.9 times 10<sup>-6</sup> M. At 10<sup>-5</sup> M or more of KB-2796, a complete suppression of the LVA I-Ca was observed in the majority of neurons tested. There was no apparent effect on the current-voltage (I-V) relationship and the current kinetics. KB-2796 delayed the reactivation and enhanced the inactivation of the Ca-2+ channel for LVA I-Ca voltage- and time-dependently, suggesting that KB-2796 preferentially binds to the inactivated Ca-2+ ~~channel~~. ~~KB~~-2796 at a concentration of 3.0 times 10<sup>-6</sup> M also decreased the peak amplitude of the HVA I-Ca without shifting the I-V relationship. In addition, KB-2796 reduced the oxidative metabolism (the formation of reactive oxygen species) of the neuron in a concentration-dependent manner with a threshold concentration of 3 times 10<sup>-6</sup> M. It is suggested that the inhibitory action of KB-2796 on the neuronal Ca-2+ influx and the oxidative metabolism, in combination with a cerebral vasodilatory action, may reduce ischemic brain damage.

? t s7/3/1-9

7/3/1

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0015137205 BIOSIS NO.: 200500043955

Regulation of CLC-Ka/barttin by the ubiquitin ligase Nedd4-2 and the serum-  
and glucocorticoid-dependent kinases

AUTHOR: Embark Hamdy M; Boehmer Christoph; Palmada Monica; Rajamanickam  
Jeyaganesh; Wyatt Amanda W; Wallisch Sabine; Capasso Giovambattista;  
Waldegger Petra; Seyberth Hannsjoerg W; \*\*\*Waldegger Siegfried\*\*\*;  
\*\*\*Lang Florian\*\*\* (Reprint)

AUTHOR ADDRESS: Inst PhysiolDept Physiol 1, Univ Tübingen, Gmelinstr 5,  
D-72076, Tübingen, Germany\*\*Germany

AUTHOR E-MAIL ADDRESS: florian.lang@uni-tuebingen.de

JOURNAL: Kidney International 66 (5): p1918-1925 November 2004 2004

MEDIUM: print

ISSN: 0085-2538 (ISSN print)

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

7/3/2

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0015038985 BIOSIS NO.: 200400409774

Serum and glucocorticoid inducible kinases functionally regulate CLC-2  
channels

AUTHOR: Palmada Monica; Dieter Michael; Boehmer Christoph; \*\*\*Waldegger\*\*\*  
\*\*\*Siegfried\*\*\*; \*\*\*Lang Florian\*\*\* (Reprint)

AUTHOR ADDRESS: Inst Physiol, Univ Tübingen, Gmelinstr 5, D-72076,  
Tübingen, Germany\*\*Germany

AUTHOR E-MAIL ADDRESS: florian.lang@uni-tuebingen.de

JOURNAL: Biochemical and Biophysical Research Communications 321 (4): p

1001-1006 September 3, 2004 2004

MEDIUM: print

ISSN: 0006-291X

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

7/3/3

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0015033115 BIOSIS NO.: 200400403904

Activating mutation of the renal epithelial \*\*\*chloride\*\*\* channel ClC-Kb  
predisposing to hypertension

AUTHOR: Jeck Nikola; \*\*\*Waldegger Siegfried\*\*\*; Lampert Angelika; Boehmer  
Christoph; Waldegger Petra; Lang Philipp A; Wissinger Bernd; Friedrich  
Bjoern; Risler Teut; Moehle Robert; Lang Undine E; Zill Peter; Bondy  
Brigitta; Schaeffeler Elke; Asante-Poku Stephen; Seyberth Hannsjoerg;  
Schwab Matthias; \*\*\*Lang Florian\*\*\* (Reprint)

AUTHOR ADDRESS: Dept Physiol, Univ Tübingen, Gmelinstr 5, D-72076,  
Tübingen, Germany\*\*Germany

AUTHOR E-MAIL ADDRESS: florian.lang@uni-tuebingen.de

JOURNAL: Hypertension (Baltimore) 43 (6): p1175-1181 June 2004 2004

MEDIUM: print

ISSN: 0194-911X

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

7/3/4

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0014294469 BIOSIS NO.: 200300253188

Regulation of channels by the serum and glucocorticoid-inducible kinase:

Implications for transport, excitability and cell proliferation.

AUTHOR: \*\*\*Lang Florian\*\*\* (Reprint); Henke Guido; Embark Hamdy M;  
\*\*\*Waldegger Siegfried\*\*\*; Palmada Monica; Bohmer Christoph; Vallon  
Volker

AUTHOR ADDRESS: Physiologisches Institut I, Gmelinstrasse 5, D-72076,  
Tuebingen, Germany\*\*Germany

AUTHOR E-MAIL ADDRESS: florian.lang@uni-tuebingen.de

JOURNAL: Cellular Physiology and Biochemistry 13 (1): p41-50 2003 2003

MEDIUM: print

ISSN: 1015-8987

DOCUMENT TYPE: Article; Literature Review

RECORD TYPE: Abstract

LANGUAGE: English

7/3/5

DIALOG(R)File 5:Biosis Previews(R)

(c) 2005 BIOSIS. All rts. reserv.

0013015107 BIOSIS NO.: 200100186946

Cloning and characterization of SLC26A6, a novel member of the solute  
carrier 26 gene family

AUTHOR: \*\*\*Waldegger Siegfried\*\*\* (Reprint); Moschen Ivano; Ramirez Alfredo  
; Smith Richard J H; Ayadi Hammadi; \*\*\*Lang Florian\*\*\*; Kubisch Christian

AUTHOR ADDRESS: Zentrum fuer Kinderheilkunde, Universitaet Marburg,  
Deutschhausstr. 12, D-35037, Marburg, Germany\*\*Germany

JOURNAL: Genomics 72 (1): p43-50 February 15, 2001 2001

MEDIUM: print

ISSN: 0888-7543

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

7/3/6

DIALOG(R)File 5:Biosis Previews(R)

(c) 2005 BIOSIS. All rts. reserv.

0012052103 BIOSIS NO.: 199900311763

Functional characterization of the betaine/gamma-aminobutyric acid  
transporter BGT-1 expressed in Xenopus oocytes

AUTHOR: Matskevitch Ioulia; Wagner Carsten A; Stegen Carola; Broer Stefan;  
Noll Birgitta; Risler Teut; Kwon H Moo; Handler Joseph S; \*\*\*Waldegger\*\*\*  
\*\*\*Siegfried\*\*\*; Busch Andreas E; \*\*\*Lang Florian\*\*\* (Reprint

AUTHOR ADDRESS: Institute of Physiology I, University of Tuebingen,  
Gmelinstrasse 5, 72076, Tuebingen, Germany\*\*Germany

JOURNAL: Journal of Biological Chemistry 274 (24): p16709-16716 June 11,  
1999 1999

MEDIUM: print

ISSN: 0021-9258

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

7/3/7

DIALOG(R)File 5:Biosis Previews(R)

(c) 2005 BIOSIS. All rts. reserv.

0010414706 BIOSIS NO.: 199699048766

Expression of a renal type I sodium/phosphate transporter (NaPi-1) induces  
a conductance in Xenopus oocytes permeable for organic and inorganic  
anions

AUTHOR: Busch Andreas E (Reprint); Schuster Andreas; \*\*\*Waldegger\*\*\*  
\*\*\*Siegfried\*\*\*; Wagner Carsten A; Zempel Guenther; Broer Stefan; Biber  
Juerg; Murer Heini; \*\*\*Lang Florian\*\*\*

AUTHOR ADDRESS: Inst. Physiol. I, Eberhard-Karls-Universitaet Tuebingen,  
Gmelinstrasse 5, D-72076 Tuebingen, Germany\*\*Germany

JOURNAL: Proceedings of the National Academy of Sciences of the United  
States of America 93 (11): p5347-5351 1996 1996

ISSN: 0027-8424  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

7/3/8

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0010108804 BIOSIS NO.: 199698576637

The type I phosphate (P-i) transporter is a novel anion channel  
AUTHOR: Busch Andreas E (Reprint); Schuster Andreas (Reprint); \*\*\*Waldegger\*\*\*  
\*\*\* Siegfried\*\*\* (Reprint); Wagner Carsten A (Reprint); Biber Juerg; Murer  
Heini; \*\*\*Lang Florian\*\*\* (Reprint  
AUTHOR ADDRESS: Univ. Tuebingen, Tuebingen, Germany\*\*Germany  
JOURNAL: Journal of the American Society of Nephrology 6 (3): p359 1995  
1995  
CONFERENCE/MEETING: Annual Meeting of the American Society of Nephrology  
San Diego, California, USA November 5-8, 1995; 19951105  
ISSN: 1046-6673  
DOCUMENT TYPE: Meeting; Meeting Abstract  
RECORD TYPE: Citation  
LANGUAGE: English

7/3/9

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2005 BIOSIS. All rts. reserv.

0009508649 BIOSIS NO.: 199497529934

Molecular basis of I-SK protein regulation by oxidation and chelation  
AUTHOR: Busch Andreas E (Reprint); \*\*\*Waldegger Siegfried\*\*\* (Reprint);  
Herzer Tobias (Reprint); Raber Gertraud (Reprint); Gulbins Erich  
(Reprint); Swanson Richard; Folander Kimberly; Takumi Toru; Moriyoshi  
Koki; Nakanishi Shigetada; \*\*\*Lang Florian\*\*\* (Reprint  
AUTHOR ADDRESS: Physiol. Inst. I, Univ. Tuebingen, Tuebingen, Germany\*\*  
Germany  
JOURNAL: Journal of the American Society of Nephrology 5 (3): p283 1994  
1994  
CONFERENCE/MEETING: Abstracts Submitted for the 27th Annual Meeting of the  
American Society of Nephrology Orlando, Florida, USA October 26-29, 1994;  
19941026  
ISSN: 1046-6673  
DOCUMENT TYPE: Meeting; Meeting Abstract; Meeting Poster  
RECORD TYPE: Citation  
LANGUAGE: English

? log y

07mar05 15:54:40 User217744 Session D904.3  
\$10.98 1.910 DialUnits File5  
\$18.00 9 Type(s) in Format 3  
\$14.00 7 Type(s) in Format 7  
\$32.00 16 Types  
\$42.98 Estimated cost File5  
\$2.93 TELNET  
\$45.91 Estimated cost this search  
\$45.93 Estimated total session cost 2.241 DialUnits  
Logoff: level 05.00.10a D 15:54:40